

The Role of a Hot Gas Environment on the Evolution of Galaxies

David Burstein¹, George Blumenthal²

Received _____; accepted _____

¹Department of Physics and Astronomy, Box 871504, Arizona State University, Tempe, AZ 85287-1504

²Board of Studies, Astronomy & Astrophysics, University of California, Santa Cruz, CA 96054

ABSTRACT

Most spiral galaxies are found in galaxy groups with low velocity dispersions; most E/S0 galaxies are found in galaxy groups with relatively high velocity dispersions. The mass of the hot gas we can observe in the E/S0 groups via their thermal X-ray emission is, on average, as much as the baryonic mass of the galaxies in these groups. By comparison, galaxy clusters have as much or more hot gas than stellar mass. Hot gas in S-rich groups, however, is of low enough temperature for its X-ray emission to suffer heavy absorption due to Galactic HI and related observational effects, and hence is hard to detect. We postulate that such lower temperature hot gas does exist in low velocity dispersion, S-rich groups, and explore the consequences of this assumption. For a wide range of metallicity and density, hot gas in S-rich groups can cool in far less than a Hubble time. If such gas exists and can cool, especially when interacting with HI in existing galaxies, then it can help link together a number of disparate observations, both Galactic and extragalactic, that are otherwise difficult to understand.

Subject headings: galaxies:formation — galaxies:evolution — intergalactic medium

1. Introduction

Both the Nearby Galaxies Catalog (Tully 1987) and the discovery of X-ray emitting hot gas in galaxy groups (Mulchaey, et al. 1993; Heldson & Ponman 2000; Mulchaey 2000, hereafter JSM00) have provided strong evidence that most galaxies reside in galaxy groups, not unlike our own Local Group (e.g., all of ~ 2500 galaxies in the Nearby Galaxies Catalog are within 1 Mpc of a galaxy group or cluster). Indeed, the evidence suggests that up to

90% of galaxies reside in galaxy groups, rather than in the better-studied galaxy clusters (cf. Tully 1987; Nolthenius 1993).

Galaxy groups can be divided into two types by the morphology of their giant galaxies (Mulchaey, et al. 1996a; Burstein, et al. 1997). The dividing line chosen by Mulchaey, et al. (1996b) is that any group that has more than 1/3 of its giant galaxies of types gE or S0 is called E/S0-rich; all others are called S-rich. Two-thirds of the Nolthenius (1993) nearby sample of galaxy groups are S-rich by this definition; 1/3 are E/S0-rich (Burstein, et al. 1997). The mean observed velocity dispersion in the galaxy groups classified by Nolthenius decreases by a factor of two from E/S0-rich to S-rich groups (Mulchaey, et al. 1996b). Subsequently, Zabludoff & Mulchaey (1998) and Mulchaey & Zabludoff (1998) found that while E/S0-rich groups can contain 30 or more galaxies more luminous than $M_B \sim -16$, typical S-rich groups, such as our own Local Group, contain only 4-8 galaxies this luminous.

Mulchaey, et al. (1996b) point out that the higher velocity dispersion of E/S0-rich groups corresponds to a hot gas temperature of ~ 0.9 keV, which is easily seen with ROSAT. In contrast, for S-rich groups, the predicted temperature based on their lower velocity dispersion averages 1/4 of this value, making any X-ray emission from hot gas subject to severe extinction by metals associated with HI gas in our own Galaxy (Raymond & Smith 1977).

Here we make the reasonable assumption that low velocity dispersion, S-rich galaxy groups *do* contain hot gas (in some S-rich groups such hot gas has been found; cf. JSM00; Xu & Wu, 2000). In Section 2 we show that, for a range of temperature, metallicity and density, much of the hot gas in galaxy groups can cool in far less than a Hubble time. In Section 3, we briefly discuss the consequences of the interrelationship that must exist among the dark matter distribution and hot gas in galaxy groups, and the consequences that relationship holds for the formation and evolution of galaxies in the groups. Section 4

discusses some of the varied observations of galaxies and galaxy groups that the assumptions of this Letter can help explain or predict.

2. Cooling Times for Hot Gas in Galaxy Groups

The measured temperatures of X-ray emitting gas in galaxy groups range from $T \sim 0.3 - 1.0$ keV, with a strong correlation between the velocity dispersion of the groups and their hot gas temperatures (e.g., JSM00). This lower temperature limit arises not from the absence of groups with low velocity dispersions (predominantly S-rich), but rather from the He and metal nuclei absorption (associated with Galactic HI) of X-ray emission from hot gas with $T \leq 0.3$ keV ($T \leq 3.5 \times 10^6$ K) (e.g., Raymond & Smith 1977). The fact that lower hot gas temperature means lower luminosity, and that the X-ray background is higher for $T < 0.5$ keV, also contribute to the lack of X-ray detected low velocity dispersion groups.

It is well-known that X-ray emitting gas with $T \sim 10^6 - 10^7$ K can cool efficiently under the “right” circumstances of gas density, metallicity and temperature (Nulsen, Stewart & Fabian 1984). In Figure 1 we plot the gas cooling times as a function of temperature for a range of temperature and metallicity relevant for gas in galaxy groups, using the results given by Sutherland & Dopita (1993). Cooling times are inversely proportional to the density of gas; the density ($n_H = 10^{-3} \text{ cm}^{-3}$) used for Figure 1 corresponds to the typical gas densities seen in the interiors of the E/S0-rich galaxy groups (cf. Mulchaey, et al. 1996a). JSM00 finds that most galaxy groups have hot gas with metallicities $z > 0.1$. Even for gas densities a factor of ten less than used for Figure 1, cooling times are still less than a Hubble time for $z > 0.1$, especially for the gas temperatures ($T < 3.5 \times 10^6$ K) we expect to find in S-rich groups.

3. Hot Gas and Galaxy Evolution

The existing data is consistent with the mass of hot X-ray emitting gas being comparable to the visible mass of galaxies, stars and gas (cf. discussion in JSM00). The fact that hot intragroup/intracluster gas contains metals implies that any primordial gas that was originally there was subsequently enriched from star formation. Consequently, during the history of these galaxies, their star formation processes must have produced a comparable amount of mass in hot gas released into the intragroup medium as there is mass in the stars themselves today. The corresponding production rate for hot gas in stellar evolution implies an average star formation rate not wildly different from what we see in our Galaxy today (cf. Bachiller 1996).

By creating this hot gas via stellar evolution, galaxies are putting a substantial amount of their mass into “hot storage”, where subsequent cooling and interactions with galaxies may significantly affect the galaxies’ structure and evolution. This is yet another reason (cf. the density-morphology relation for galaxies, Dressler 1980) to view the formation and evolution of galaxies not in individual terms, but rather in the context of the galaxy groups and clusters to which they belong.

The mass-to-light ratios for galaxy groups, derived from the X-ray observations, are similar to those for galaxy clusters, 100-120 in solar units, B mag (cf. JSM00), implying dark matter domination. The mean velocity dispersion for S-rich groups is 1/2 that of E/S0-rich groups (cf. above), and E/S0-rich groups also have effective radii that are 50% larger than S-rich groups (cf. data in Burstein, et al. 1997). On average, E/S0-rich groups are more massive ($6\times$) and denser ($1.7\times$) than S-rich groups. Because of its tight correlation with velocity dispersion, the hot gas temperature in galaxy groups is determined by the density and mass of the group, which in turn is dominated by dark matter.

It is also noteworthy that the velocity dispersions *within* large galaxies in groups

tend to be comparable to the velocity dispersions of the galaxies *in* their groups: E/S0 galaxies have typical velocity dispersions (at L^*) of 250 km s^{-1} , while giant spirals typically have equivalent velocity dispersions of $150\text{-}200 \text{ km s}^{-1}$ (Burstein, et al. 1997). Obviously, this trend breaks down at the higher masses of clusters with their much higher velocity dispersion, but there is evidence that clusters represent the end point of the clustering of groups in dense environments (Blumenthal et al. 1984). Within galaxy groups, the “dark matter/visible matter conspiracy” corresponding to the similarity of stellar motions and dark matter velocity dispersion therefore applies to both the environment of galaxies and to the galaxies themselves (cf. Burstein & Rubin 1985).

This view is consistent with the hierarchical-clustering-merging (HCM) picture (Burstein, et al. 1997) in which E/S0 galaxies are formed in denser, more massive environments, while spirals are formed in lower mass, less dense environments. The new wrinkle here is something that the HCM scenario has trouble predicting quantitatively (owing to a lack of physical understanding of the origin of the initial mass function for stars). Namely, galaxies put much of their baryonic mass into “hot” storage. Some of this hot gas will be in storage in the halo of the galaxies themselves, while some will be within the general group medium.

The fact that much of this hot gas can cool efficiently to cold gas in less than a Hubble time can have important observational implications. The interaction of hot intragroup gas with the cold gas in galaxies moving within the groups can lead to complex hydrodynamical effects including shocks and phase transitions, with subsequent cooling. This interaction of hot and cold gas can lead to several interesting galactic phenomena (e.g., tidal tails, galactic fountains, small, HI-rich dwarf galaxies in orbit about their parent galaxies, such as in the Local Group). In fact, shocks that form in the interaction region lead to higher densities for the hot gas, with consequently shorter cooling times. Quantitatively addressing these

complications will involve detailed modeling with yet-to-be-written N-body hydro codes, but it is reasonable to assume that the cooling of this hot gas and its interactions with extant galaxies can significantly affect the evolution of these galaxies. We now explore some of the plausible observational consequences of this interaction.

4. Observational Consequences of Galaxy Interactions with Intragroup Gas

For the following, we make three assumptions: a) hot gas exists in all galaxy groups; b) much of this gas can cool in much less than a Hubble time; and c) the cooling rate is even faster when there are shocks associated with the interstellar medium in galaxies. These assumptions lead to a wide range of linked observational implications.

1. The density/morphology relation for galaxies: It has long been known (e.g. Dressler 1980) that galaxy morphology correlates with local initial conditions such as mass density. Almost certainly, part of this correlation is causal, since higher density implies more major merging, which probably leads to ellipticals. At the same time, higher initial densities lead to groups with higher velocity dispersion and gas temperature. While the gas in S-rich groups will therefore be able to cool in less than a Hubble time, the hotter gas E/S0-rich groups may be too hot ($\sim 10^7$ K, cf. JSM00 and Fig. 1) to efficiently cool.

2. Galaxy warps, tidal tails and HI mass surface density distributions: Most edge-on spiral disks have warps that persist well into their extended HI distribution (Sancisi 1976), which is consistent with the warp model developed to match the HI distribution (Bosma 1981). Surprisingly, there is evidence that the ratio of mass surface density (mostly dark matter) to HI surface density in the outer regions of spirals is a constant over several scale lengths (Bosma 1981; Hoekstra, Sancisi, & van Albada 2001), with a sharp outer cut-off in the HI surface density (Hoekstra et al.). Some HI-dominated tidal tails extending

from spiral galaxies have more gas than likely just from the galaxies from which they are extended (Hibbard & Yun 2001).

These observations of spiral galaxies are consistent with a strong continuing interplay between cold HI in galactic disks and a cooling, hot intragroup medium. In particular, hot gas may cool to form a part of the HI disk, or there may be a pressure equilibrium between the colder disk material and the hot gas. Tidally-extruded HI from galaxies can possibly stimulate formation of HI from the hot gas that surrounds the galaxies, likely forming the HI in pressure equilibrium with the hot gas. Only detailed hydro modeling can tell us more about how this interaction might work, but the *prima facie* evidence is that there is simply too much mass in HI tidal tails to have come just from the outer, low HI surface density parts of spirals. The observed sharp cutoffs in the HI disks of galaxies might also arise from dynamical pressure or cooling from the hot gas surrounding the galaxy.

3. High velocity HI clouds: These exist near our Galaxy, with their distances and origin still hotly debated (Blitz et al. 1997; Sembach 2002a,b). Recent observations (cf. Sembach 2002a,b) indicate that the high velocity clouds near our Galaxy are enveloped in hot gas of temperature $\sim 10^6$ K. Similar kinds of “high velocity clouds” are observed near the disks of other spiral galaxies (Sancisi, et al. 2001), suggesting that this phenomenon is not isolated. It is possible that these cold clouds could arise from thermal cooling stimulated by the passage of dwarf galaxies or by an outflowing wind driven by star formation in galaxy disks. Similarly, when we look at the Galactic poles in both directions, we see HI gas infalling toward the Galactic plane (cf. Tolbert 1971). This is consistent with the idea that a local blowout for a galaxy disk can stimulate cold gas to form from the hot gas in the galaxy’s halo, and fall back to the Galactic plane.

4. Ultraviolet spectroscopic observations of high ionization absorption features in distant objects show the presence of hot gas near our Galaxy along different lines of sight

(e.g., Dixon, Davidsen, & Ferguson 1995; Savage et al. 2002; Sembach 2002b) as well as in other galaxy groups (e.g., Tripp 2002). As pointed out by Mulchaey, et al. (1996b), this is precisely what hot gas in S-rich groups (or in our galactic halo and/or in our Local Group) would produce.

5. Galaxy groups which show strong tidal interactions among their members often have a substantial fraction of their HI outside their galaxies. For example, the M81 group has 25% of its HI gas in its intragroup medium, rather than in the galaxies (Appleton, Davies, & Stepenson 1981). This does not include the HI gas from the intragroup medium that has already been dumped on M82 and NGC 3077 and formed stars there. It is unlikely that tidal tails alone can produce such a high percentage of HI gas in the general group medium (again, they arise from the relatively low density outer parts of galaxies). Much of the intragroup gas in the M81 group could have come from cooling of the hot gas in that group, stimulated by the tidal interactions of the galaxies, as well as with subsequent star formation in M82 and NGC 3077.

6. The solution of the “G-dwarf problem” in our Galaxy most likely requires a constant infall of low metallicity gas (cf. Chiappina, Matteucci, & Gratton 1997). Similarly, semi-analytic models of galaxy evolution seem to imply the need for continuous infall of gas from outside galaxies to produce the galaxies we see today (cf. Sembach 2002a). Hot gas in the halo or the immediate environment of spiral galaxies, which is of moderately low metallicity, and which can cool to cold gas, can provide just such a reservoir.

7. Spiral arms appear in HI disks of spirals outside the disks of stars (e.g., Bosma 1981). Given that galaxies move at roughly the speed of sound through their intragroup gas, the shocks that are produced can help stimulate spiral arm formation in the HI gas in the outer parts of galaxies.

8. The relationship between kinematic velocity fields and HI distributions in spiral

galaxies is complicated. While the kinematic velocity fields in many non-interacting spiral galaxies are typically very regular, the angular distribution of HI gas (and, to some extent, the disk stars) can be very irregular (e.g., Bosma 1981). In extreme cases, the HI density distribution within a spiral galaxy disk can be very lopsided, while preserving a regular velocity field. There are exceptional cases in which the kinematic velocity fields can also be lopsided (Swaters, et al. 1999). If cold gas is being produced by cooling of hotter gas due to interactions with, for example, gas produced by star formation in the galaxy disks, this could easily lead to a highly variable gas density without significantly affecting the kinematic velocity field. This also suggests that the disks form from the inside out (higher gas density in the galaxy interiors), producing blue color gradients owing both to lower metallicity and younger stars as a function of increasing radius (cf. Courteau & Holtzman 1995).

Of course, the above is not an exhaustive list of observational consequences of our assumptions. Moreover, not every one of these observations is necessarily a consequence of our assumptions. Rather, we suggest that all of the observations listed here are *consistent* with our assumptions. Based on this discussion, we have estions for future investigation:

- i. Density influences the formation and evolution of galaxies in more than just defining the circumstances of their birth. Rather, the temperature of the hot gas they produce in their groups also can dictate their subsequent evolution. Hydro modeling of galaxies must take into account the environments in which we find them, which are group-dominated.
- ii. Observations indicate a range in gas mass in galaxy groups (JSM00), some of which may be observationally-driven, but some of which may be real. Separately, spiral galaxies are found with a wide range in surface brightness (cf. Impey et al. 1996). There could be a relationship between gas mass/gas temperature ranges in galaxy groups and the surface brightness of their spiral galaxies that should be explored.

iii. Similarly, HI content and properties of spiral galaxies would depend on the galaxy groups in which we find them. The known large variation of HI properties among spiral galaxies (Bosma 1981; Hoekstra, Sancisi, & van Albada 2001) makes this prediction only testable with a large statistical sample of galaxies in a wide range of group environments. Unfortunately, most of the nearby, large spirals which have been well-studied for their HI distribution are part of the Coma-Sculptor Cloud, which is completely comprised of S-rich groups (Tully 1987).

iv. Of course, direct detection of hot gas in low dispersion galaxy groups would directly prove our assumptions. Unfortunately, the very long exposure times necessary for this to happen make such observations hard to do.

If the way galaxy formation and evolution proceeds as we have outlined in this Letter is correct, then two key pieces of information are missing from high- z galaxy images: the three-dimensional distribution of dark matter and the kinds of hot gas that are found at each stage of galaxy formation and evolution. Knowing the temperature and density of hot gas in groups as a function of redshift would allow one to make better observational predictions for different ranges of time.

In closing, it is clear that many of the possible observational consequences we have identified can be modeled numerically. In principle, N-body hydro codes can adequately model the interactions of hot gas in galactic groups with the disks of spiral galaxies. However, the star formation process and the way that star formation relates to galaxy (and surrounding group) properties may turn out to be an essential element. Until we understand the physics that produces the initial mass functions of stars, any hypotheses such as ours that involve interaction of hot gas in galaxy groups with cold gas and star formation in galaxies must remain at least partly speculative.

We wish to thank Dr. George Coyne, SJ, Dr. Jose’ Funes, SJ, Dr. Enrico Corsini and Professor Francesco Bertola for organizing the June 2000 meeting on “Galaxy Disks and Disks in Galaxies”, and conversations with Renzo Sancisi, which greatly stimulated our ideas.

REFERENCES

- Appleton, P.N, Davies, R.D., & Stepenson, R.J., 1981, MNRAS, 195, 327
- Bachiller, R. 1996, ARA&A, 34, 111
- Blitz, L, Spergel, D.N., Teuben, P.J., Hartmann, L., & Burton, W.B., 1997, BAAS, 28, 1349
- Blumenthal, G., Faber, S. Primack, J. & Rees, M., 1984, Nature, 311, 517
- Bosma, A., 1981, AJ, 86, 1825
- Burstein, D., Bender, R., Faber, S.M., & Nolthenius, R., 1997, AJ, 114, 1365
- Burstein, D. & Rubin, V.C., 1985, ApJ, 297, 423
- Chiappina, C., Matteucci, F., & Gratton, R., 1997, ApJ, 477, 765
- Courteau, S. & Holtzman, J. 1995, in The Opacity of Spiral Disks, eds. Davies, J.I. & Burstein, D., Kluwer, Netherlands, 211
- Dixon, W.V., Davidsen, A.F., & Ferguson, H.C., 1995, BAAS, 27, 1452
- Dressler, A., 1980, ApJ, 236, 351
- Dressler, A. & Shectman, S.A., 1988, AJ, 95, 985
- Heldson, S.F. & Ponman, T.J., 2000, MNRAS, 315, 356
- Hibbard, J. & Yun, M.S., 2001, private communication, see figure at
www.cv.nrao.edu/~hibbard/LIRpaper/LIR2_fig4.gif
- Hoekstra, H., Sancisi, R., & Van Albada, T.S., 2001, MNRAS, 323, 453
- Impey, C. D., Sprayberry, D., Irwin, M. J., & Bothun, G. D. 1996, ApJS, 105, 209

- Mulchaey, J.S., 2000, ARA&A, 38, 289
- Mulchaey, J.S., Davis, D.S., Mushotzky, R.F., & Burstein, D., 1993, ApJ, 404, 9
- Mulchaey, J.S., Davis, D.S., Mushotzky, R.F., & Burstein, D., 1996a, ApJ, 456, 80
- Mulchaey, J.S., Davis, D.S., Mushotzky, R.F., & Burstein, D., 1996b, ApJ, 456, 5
- Mulchaey, J.S. & Zabludoff, A.I., 1998, ApJ, 496, 73
- Nolthenius, R., 1993, ApJS, 85, 1
- Nulsen, P.E.J., Stewart, G.C., & Fabian, A.C., 1984, MNRAS, 208, 185
- Raymond, J.C. & Smith, B.W., 1977, ApJS, 419
- Sancisi, R., 1976, A&A, 53, 159
- Sancisi, R., Fraternali, F., Oosterloo, T., & van Moorsel, G., 2001, in Galaxy Disks and Disk Galaxies, ASP Conf. Ser. 230, edited by J.G. Funes, S.J. & E.M. Corsini, (ASP, San Francisco), p. 111
- Savage, B.D., Sembach, K.R., Wakker, B.P., Richter, P., & Meade, M., 2002, BAAS, 33, 1407
- Sembach, K.R., 2002, BAAS, 33, 1478
- Sembach, K.R., 2002, Space Tel Sci Newsletter, 19, 2, 19 (Spring 2002)
- Sutherland, R.S. & Dopita, M.A., 1993, ApJS, 88, 253
- Swaters, R.A., Schoenmakers, R.H..M., Sancisi, R., & van Albada, T.S., 1999, MNRAS, 304, 330
- Tolbert, C.R., 1971, A&AS, 3, 349

Tripp, T.M, 2002, BAAS, 34, 689

Tully, R.B., 1987, Nearby Galaxies Catalog, Cambridge University Press, New York, N.Y.

Xue, Y.-J. & Wu, X.-P., 2000, ApJ, 538, 65

Zabludoff, A.I. & Mulchaey, J.S., 1998, ApJ, 496, 39




Fig. 1.— The logarithm of the cooling rates (in years) of hot gas as a function of the logarithm of its temperature (given both in terms of K and in keV), for four values of metallicity relative to solar (z); 1.0, 0.1, 0.01 and 0.00. Dotted lines are drawn between the $z = 1.0$ and $z = 0.1$ curves at values of $\text{Log } T = 5.78$ and 6.5 , corresponding to the range of observed velocity dispersions in low- σ galaxy groups, for a gas density of $n_{\text{H}} = 10^{-3} \text{ cm}^{-3}$. A dotted line at a cooling time of 1 Gyr is drawn to guide the eye.